

SPECTROPHOTOMETRIC DETERMINATION OF SECNIDAZOLE USING FOLIN CIOCALTEU'S & SODIUM CARBONATE

K. Senthil kumar¹, B. Manasa¹, E. Nagamani¹, G. Manoj Varma¹, Errolla Mahesh¹, B.Sudhakar² and TV. Narayana³

¹Sri Adichunchanagiri College of Pharmacy, B.G nagara, Karnataka, Bangalore, India.

²The Himalaya drug company, Bangalore, Karnataka, India.

³Chairman, Dr.H.L.T College of Pharmacy,Channapatna,Ramanagara Dist Karnataka,India.

ABSTRACT

Absorption spectroscopic methods based on ultraviolet and visible radiation is widely used for the determination of a variety of organic and inorganic compounds. The proposed method is simple,sensitive,accurate and precise.secnidazole treated with folin-ciocalteu reagent and sodiumcarbonate to form blue coloured chromogen. The absorbance of the solution was measured at 677nm. Beers law was obeyed in the concentration range of 5-25mcg/ml.

The optical charateristics such as absorption such as absorption maxima,beers law limit, correlation co-efficient,slope,intercept, 1regression equation,relative standard deviation and recovery studies were also carried out. Results were validated statistically and were found to be reproducible. The analysis results of marketed formulation compiles with their labelled claim.

Keywords: Secnidazole, Spectrophotometry, Folin Ciocalteu, Sodium Carbonate.

PREPARATION OF STANDARD STOCK SOLUTION

100 mg of secnidazole was accurately weighed and dissolved in 30ml of distilled water and treated with 10ml of 4N hydrochloric acid and 1.2gm of zinc dust was added in portions. After standing for 1hr at room temperature the solution was filtered through cotton wool, the residue was washed with 3×10ml portions of distilled water and the total volume of the filtrate was brought to 100ml with distilled water (solution I)

ABSORPTION SPECTRA OF COLOURED SPECIES

25ml of solution I was pipette into 100ml standard volumetric flask and was diluted to 100ml with distilled water. (solution II)


From the solution II 2ml was pipetted out into a dry 25ml standard volumetric flask. 2.5ml of folin ciocalteu's reagent, and 5ml of sodium carbonate solution were added and the volume was made upto 25ml with sodium hydroxide solution. The absorption spectrum was obtained by scanning between the wavelength 550-740 nm against the reagent blank.

The readings are tabulated in table no:1a and graphically shown in graph no: 1a

Table 1a: Absorption Spectral Data of Coloured Species Using Folin Ciocalteu's & Sodium Carbonate Reagent
Drug concentration 20 μ g/ml

Wavelength in nm	Absorbance
550	0.231
560	0.235
570	0.234
580	0.236
590	0.237
600	0.238
610	0.241
620	0.243
630	0.244
640	0.246
650	0.247
660	0.248
670	0.250
680	0.251
690	0.247
700	0.245
710	0.240
720	0.237
730	0.233
740	0.232

Graph 1a: Absorption Spectra for Drug With Folin Ciocalteu's and Sodium Carbonate Drug concentration- 20 μ g/ml



e. EFFECT OF REAGENT CONCENTRATION

To find out the effect of different amount of folin ciocalteu's reagent 1ml, 1.5ml, 2.0ml, 2.5ml of folin ciocalteu's reagent was added into 5, 10, 15, 20 & 25 μ g/ml concentration of drug. 5ml of sodium carbonate solution was


also added. The absorbance was measured at 677nm against the reagent blank.

The readings are shown in table no 1.b and are graphically represented in graph no 1.b, 1.c, 1.d, 1.e


Table 1b: Data for the Calibration Curve With Different Volume of Folin Ciocalteu's Reagent

S.NO	Drug concentration in $\mu\text{g/ml}$	Volume of sodium carbonate reagent added	Volume of folin ciocalteu's reagent added & Absorbance			
			1ml	1.5ml	2ml	2.5ml
1	5	5ml	0.057	0.059	0.061	0.062
2	10	5ml	0.119	0.121	0.122	0.124
3	15	5ml	0.179	0.180	0.181	0.185
4	20	5ml	0.231	0.235	0.241	0.244
5	25	5ml	0.289	0.292	0.298	0.302


**Graph 1b: Calibration Curve for Various Concentration of Drug With 1 ml of Folin Ciocalteu's Reagent & 5 ml of Sodium Carbonate Reagent
ABSORBANCE AT 677 nm**




**Graph 1C: Calibration Curve For Various Concentration of Drug With 1.5ml of Folin Ciocalteu's Reagent & 5ml of Sodium Carbonate Reagent
ABSORBANCE AT 677 nm**



Graph 1d: Calibration Curve for Various Concentration of Drug With 2ml of Folin Ciocalteu's Reagent & 5ml of Sodium Carbonate Reagent



Graph 1e: Calibration Curve for Various Concentration of Drug with 2.5ml of Folin Ciocalteu's Reagent & 5ml of Sodium Carbonate Reagent.



f. FIXATION OF VARIOUS PARAMETERS

λ_{max} (maximum wavelength)

The spectral data shows that the maximum absorbance was observed at 677nm and that was selected as working maximum.

2. Stability of colour

Stability of the colour of this system was tested by using 5, 10, 15, 20 & 25 mcg/ml

concentration of drug with 2.5ml of folin ciocalteu's reagent and 5ml of sodium carbonate solution. Absorbance was measured at 677nm against the reagent blank. The readings were taken at 15min time interval upto 1hr. The results are shown in table 1.d

Table 1d: Data For Stability of Colour

Concentration of drug in mcg/ml	Time interval & Absorbance at 677nm				
	0 mins	15 mins	30 mins	45 mins	60 mins
5	0.062	0.062	0.062	0.062	0.059
10	0.122	0.122	0.122	0.122	0.124
15	0.181	0.181	0.181	0.181	0.178
20	0.242	0.242	0.242	0.242	0.244
25	0.303	0.303	0.303	0.303	0.298

3. Specific Extinction co-efficient

This was found by
 $= A/Lc = 125.5$

4. Beer's law plot

Beer's law plot was constructed by measuring the absorbance of various concentration of drug solution against the reagent blank.


From solution II, 1ml, 1.5ml, 2ml, 2.5ml & 3ml were taken into five different dry 25ml

standard volumetric flasks. To that 2.5ml of folin ciocalteu's reagent and 5ml of sodium carbonate were added respectively. The colour was allowed to develop, the volume was made up to the mark by slowly adding sodium hydroxide solution. The absorbance was measured at 677nm against the reagent blank. The readings are shown in table no 1.e graphically plotted in graph no 1.f

Table 1e: Data for Beer's Law Plot

Concentration of drug in mcg/ml	Absorbance at 677 nm
5	0.063
10	0.122
15	0.182
20	0.243
25	0.304

Graph 1f: Beer's Law Plot for Secnidazole With Folin Ciocalteu's And Sodium Carbonate At 677nm



g. Assay of tablet

Three different brands of secnidazole (Ambiform, Seczol DS, Seclong DS) were taken for analysis.



Total weight of each brand of tablet was determined. Powdered tablet equivalent to 100mg was accurately weighed and dissolved in 30ml of distilled water and treated with 10ml of 4N hydrochloric acid and 1.2 gm of zinc dust was added in portions. After standing for 1hr at room temperature the solution was filtered through cotton wool. The residue was washed with 3x10 ml portions of distilled water and the total volume of the filtrate was brought to 100ml with distilled water (solution I)

From solution I, 25ml was pipetted out into a dry 100ml standard volumetric flask and the volume was made upto mark with distilled water (Solution II)

From solution II, 1.5ml was pipetted out into a dry 25ml standard volumetric flask. To that 2.5ml of folin ciocalteu's reagent and 5ml of sodium carbonate were added and the volume made upto mark with sodium hydroxide solution. The absorbance was measured at 677nm against the reagent blank.

The results are shown in table no: 1.e

The same procedure was repeated five times for each brand of tablet powder.

STANDARD DILUTION**TEST DILUTION**

*2.5ml Folin Ciocalteu's reagent & 5ml Sodium Carbonate reagent

The contents of drug was determined by using the formula

$$= \frac{\text{Test absorbance}}{\text{std. absorbance}} \times \frac{\text{std.wt}}{100} \times \frac{25}{100} \times \frac{1.5}{25} \times \frac{100}{\text{test wt}} \times \frac{100}{25} \times \frac{25}{1.5} \times \text{Total wt. of tablet}$$

CONCLUSION

The method found to be simple, economical, selective and more sensitive than most of the spectrophotometric reported methods. The statistical parameters and recovery study data clearly indicate the reproducibility and accuracy of the method. This approach could be considered for the determination of the cited drugs in the quality control laboratories.

REFERENCES

1. Sekhar K Ravi, Naidu MUR, Sekhar EC, Rao TRK, Shobha JC, Rani PU and Kumar J. Surya. Rapid and selective analysis of secnidazole in human plasma using high-performance liquid chromatography with ultraviolet detection, *J Chromatography B: Biomedical Sciences and Applications*, 1997;691(1):208-211.
2. Qi, -Y, Shan HJ and WuS. Reverse-phase HPLC determination of secnidazole and its related substances. *J Zhongguo Xinyao Zazhi*. 2002;11(4):294-296.
3. Radi AE and Hassanein A. Determination of secnidazole in urine by adsorptive stripping voltammetry. *J Chem-Pharm-Bull*. 2002;48(5):600-602.
4. Saffaj M, Charrouf A, Abourriche Y, Aboud A, Bennamara and Berrada M. Spectrophotometric determination of

- Metronidazole and Secnidazole in pharmaceutical preparations based on the formation of dyes, J Berrada Dyes and pigments. 2006;70:259-262.
5. Gandhi R, Pillai O, Thilagavathi R, Gopalakrishnan B, Kaul CL and Panchagnula R. Eur J Pharm Sci. 2002;16:175-184.